

Maxima and minima: critical points, the closed-interval method and optimization problems

Ten single-variable extremum problems: a cubic classified by the second-derivative test, a flat point that is not an extremum, a minimum the second-derivative test cannot see, the closed-interval method with an endpoint maximum, a cusp where the derivative is undefined, $x e^{-x}$, an open box, the closest point on a curve, a can of least surface area, and the mean as the minimizer of squared error, with worked solutions and the mistakes of forgetting endpoints and cusps and of reading $f' = 0$ as a guaranteed extremum.

BEFORE YOU START

Training a model is minimizing a loss, and in one variable the whole theory fits on a page: an interior minimum or maximum can only sit where the derivative is zero or undefined, and on a closed interval the endpoints are candidates too. The derivative test then sorts the candidates, and comparing values picks the winner. Problem 10 is the one-variable version of least squares: the constant that minimizes squared error is the mean. The work goes wrong in a few places: endpoints left off the list, a zero derivative taken as proof of an extremum, an inconclusive second-derivative test read as "no extremum", a point where the derivative is undefined never checked, and the location of the maximum reported instead of its value. These ten problems cover each of them.

- A **critical point** of f is an interior point where $f'(x) = 0$ or $f'(x)$ does not exist. Every local maximum or minimum inside the domain is at a critical point, but not every critical point is one.
- **First-derivative test:** if f' changes from $+$ to $-$ at a critical point, it is a local maximum; from $-$ to $+$, a local minimum; if f' keeps the same sign on both sides, it is neither.
- **Second-derivative test:** at a point with $f'(c) = 0$, $f''(c) < 0$ means a local maximum and $f''(c) > 0$ a local minimum. If $f''(c) = 0$ the test says nothing, and the first-derivative test decides.
- **Closed-interval method:** a continuous f on $[a, b]$ has a largest and a smallest value. Evaluate f at every critical point in (a, b) and at a and b ; the largest value is the maximum and the smallest the minimum.
- **Optimization problems:** name the quantity to optimize, write it as a function of one variable using the constraint, find that variable's allowed range, and then use the methods above. Answer the question that was asked: a location, a value, or both.
- $x^{1/3}$ is the real cube root, defined for negative x too, so $x^{2/3} = (x^{1/3})^2 \geq 0$ for every x .

Builds on: Differentiation rules: chain, product, quotient

Problems

1. Find the local maxima and minima of $f(x) = x^3 - 6x^2 + 9x + 1$ and classify them with the second-derivative test. difficulty ·

2. Find and classify the critical points of $f(x) = x^4 - 4x^3$. difficulty ··

3. Find the minimum of $f(x) = x^4 - 4x^3 + 6x^2 - 4x$. difficulty ··

4. Find the maximum and minimum values of $f(x) = x^3 - 3x$ on $[-2, 3]$. difficulty ·

5. Find the maximum and minimum values of $f(x) = x^{2/3}(5 - x)$ on $[-1, 4]$. difficulty ···

6. Find the maximum value of $f(x) = xe^{-x}$ for $x \geq 0$. difficulty ··

7. An open-top box is made from a 12 cm by 12 cm sheet by cutting an x by x square from each corner and folding up the sides. Which x gives the largest volume, and what is that volume? difficulty ··

8. Find the point on the curve $y = \sqrt{x}$ closest to $(4, 0)$, and the distance. difficulty ···

9. A closed cylindrical can must hold 16π cm³. Which radius and height use the least metal (the least total surface area), and what is that area? difficulty ··

10. A model predicts the same number m for the four targets 1, 2, 6 and 7. Which m minimizes the squared error $E(m) = \sum_i (m - a_i)^2$, and what is the minimum error? difficulty ··

Answers

1. Local maximum 5 at $x = 1$; local minimum 1 at $x = 3$
2. $x = 3$ is a local (and global) minimum, $f(3) = -27$; $x = 0$ is neither a maximum nor a minimum
3. Minimum -1 at $x = 1$
4. Maximum 18 at $x = 3$; minimum -2 , reached at both $x = -2$ and $x = 1$
5. Maximum 6 at $x = -1$; minimum 0 at $x = 0$
6. Maximum value $\frac{1}{e} \approx 0.368$, at $x = 1$
7. $x = 2$ cm, volume 128 cm^3
8. The point $(\frac{7}{2}, \sqrt{\frac{7}{2}}) \approx (3.5, 1.871)$, at distance $\frac{\sqrt{15}}{2} \approx 1.936$
9. $r = 2$ cm, $h = 4$ cm, area $24\pi \approx 75.40 \text{ cm}^2$
10. $m = 4$, the mean; minimum squared error 26

Worked solutions

Problem 1

Find the local maxima and minima of $f(x) = x^3 - 6x^2 + 9x + 1$ and classify them with the second-derivative test.

$f'(x) = 3x^2 - 12x + 9 = 3(x - 1)(x - 3)$	Differentiate and factor out the 3.
Critical points $x = 1$ and $x = 3$	f' is a polynomial, defined everywhere, so the only critical points are its zeros.
$f''(x) = 6x - 12$, so $f''(1) = -6$ and $f''(3) = 6$	Second-derivative test.
$f(1) = 1 - 6 + 9 + 1 = 5$ and $f(3) = 27 - 54 + 27 + 1 = 1$	The values at the critical points.
Local maximum 5 at $x = 1$; local minimum 1 at $x = 3$	$f''(1) < 0$ curves down and $f''(3) > 0$ curves up. Neither is a global extremum: a cubic goes to $\pm\infty$.

Problem 2

Find and classify the critical points of $f(x) = x^4 - 4x^3$.

$f'(x) = 4x^3 - 12x^2 = 4x^2(x - 3)$	Factor out $4x^2$.
Critical points $x = 0$ and $x = 3$	The zeros of f' .
$f''(x) = 12x^2 - 24x$, so $f''(3) = 108 - 72 = 36 > 0$	$x = 3$ is a local minimum, with $f(3) = 81 - 108 = -27$.
$f''(0) = 0$	The second-derivative test is inconclusive at 0; use the first-derivative test.

$f'(-1) = 4 \cdot 1 \cdot (-4) = -16 < 0$ and $f'(1) = 4 \cdot 1 \cdot (-2) = -8 < 0$	The factor $4x^2$ is never negative, so the sign of f' near 0 is the sign of $x - 3$, negative on both sides.
$x = 3$ is a local (and global) minimum, $f(3) = -27$; $x = 0$ is neither a maximum nor a minimum	f is decreasing straight through $x = 0$: it has a flat tangent there, not a turn. The minimum at 3 is global because $f \rightarrow +\infty$ in both directions and 3 is the only turning point.

Problem 3

Find the minimum of $f(x) = x^4 - 4x^3 + 6x^2 - 4x$.

$f'(x) = 4x^3 - 12x^2 + 12x - 4 = 4(x - 1)^3$	The coefficients 1, -3, 3, -1 are those of $(x - 1)^3$.
Critical point $x = 1$ only	$(x - 1)^3 = 0$ only at $x = 1$.
$f''(x) = 12(x - 1)^2$, so $f''(1) = 0$	Inconclusive: this does not mean there is no extremum.
$f'(x) < 0$ for $x < 1$ and $f'(x) > 0$ for $x > 1$	A cube has the sign of its base. f' changes from - to +, so $x = 1$ is a local minimum by the first-derivative test.
$f(1) = 1 - 4 + 6 - 4 = -1$	In fact $f(x) = (x - 1)^4 - 1$, which shows the same thing directly.
Minimum -1 at $x = 1$	It is global: f decreases on $(-\infty, 1)$ and increases on $(1, \infty)$.

Problem 4

Find the maximum and minimum values of $f(x) = x^3 - 3x$ on $[-2, 3]$.

$f'(x) = 3x^2 - 3 = 3(x - 1)(x + 1)$	Critical points $x = -1$ and $x = 1$, both inside $(-2, 3)$.
--------------------------------------	--

$f(-1) = -1 + 3 = 2$ and $f(1) = 1 - 3 = -2$	Values at the critical points.
$f(-2) = -8 + 6 = -2$ and $f(3) = 27 - 9 = 18$	Values at the endpoints.
Maximum 18 at $x = 3$; minimum -2, reached at both $x = -2$ and $x = 1$	Compare all four values. The maximum is at an endpoint, where $f'(3) = 24 \neq 0$: a derivative test alone would never find it.

Problem 5

Find the maximum and minimum values of $f(x) = x^{2/3}(5 - x)$ on $[-1, 4]$.

$f(x) = 5x^{2/3} - x^{5/3}$	Multiply out so that each term is a power.
$f'(x) = \frac{10}{3}x^{-1/3} - \frac{5}{3}x^{2/3} = \frac{5}{3}x^{-1/3}(2 - x) = \frac{5(2 - x)}{3x^{1/3}}$	Factor out $\frac{5}{3}x^{-1/3}$; then $x^{2/3} = x^{-1/3} \cdot x$.
$f'(2) = 0$, and $f'(0)$ is undefined	$x = 0$ makes the denominator 0; f itself is continuous there, with a cusp. Both are critical points.
$f(0) = 0$ and $f(2) = 2^{2/3} \cdot 3 \approx 4.762$	Values at the critical points.
$f(-1) = (-1)^{2/3} \cdot 6 = 6$ and $f(4) = 4^{2/3} \cdot 1 \approx 2.520$	Values at the endpoints; $(-1)^{2/3} = ((-1)^{1/3})^2 = 1$.
Maximum 6 at $x = -1$; minimum 0 at $x = 0$	The minimum is at the cusp, where f' does not exist, and the maximum is at an endpoint. The only point with $f' = 0$, $x = 2$, is a local maximum but not the global one.

Problem 6

Find the maximum value of $f(x) = xe^{-x}$ for $x \geq 0$.

$f'(x) = e^{-x} - xe^{-x} = (1 - x)e^{-x}$	Product rule; $(e^{-x})' = -e^{-x}$.
Critical point $x = 1$	$e^{-x} > 0$, so $f' = 0$ only where $1 - x = 0$.
$f' > 0$ on $[0, 1)$ and $f' < 0$ on $(1, \infty)$	f rises and then falls, so $x = 1$ is the global maximum on $[0, \infty)$.
$f(1) = e^{-1}$	Evaluate f at the maximizer.
Maximum value $\frac{1}{e} \approx 0.368$, at $x = 1$	$f(0) = 0$ and $f \rightarrow 0$ as $x \rightarrow \infty$: the exponential beats the linear factor.

Problem 7

An open-top box is made from a 12 cm by 12 cm sheet by cutting an x by x square from each corner and folding up the sides. Which x gives the largest volume, and what is that volume?

$V(x) = x(12 - 2x)^2$	The base is $12 - 2x$ on each side and the height is x .
$0 \leq x \leq 6$	The cut cannot be negative or use more than half the side; $V(0) = V(6) = 0$.
$V'(x) = (12 - 2x)^2 - 4x(12 - 2x) = (12 - 2x)(12 - 6x)$	Product rule, with $\frac{d}{dx}(12 - 2x)^2 = -4(12 - 2x)$; factor out $12 - 2x$.
Critical point $x = 2$ inside $(0, 6)$	$x = 6$, the other zero, is an endpoint.
$V(2) = 2 \cdot 8^2 = 128$	Larger than the endpoint values 0.
$x = 2$ cm, volume 128 cm^3	The box is 8 by 8 by 2.

Problem 8

Find the point on the curve $y = \sqrt{x}$ closest to $(4, 0)$, and the distance.

$D(x) = (x - 4)^2 + (\sqrt{x})^2 = (x - 4)^2 + x$, for $x \geq 0$	The squared distance from $(x, \sqrt{x})$ to $(4, 0)$. The square root is increasing, so the distance and its square have the same minimizer, and the square has no root to differentiate.
$D'(x) = 2(x - 4) + 1 = 2x - 7$	Differentiate.
Critical point $x = \frac{7}{2}$; $D'' = 2 > 0$	A minimum, and global since D is a parabola opening up. The endpoint gives $D(0) = 16$, larger.
$D(\frac{7}{2}) = \frac{1}{4} + \frac{7}{2} = \frac{15}{4}$	$(\frac{7}{2} - 4)^2 = \frac{1}{4}$.
The point $(\frac{7}{2}, \sqrt{\frac{7}{2}}) \approx (3.5, 1.871)$, at distance $\frac{\sqrt{15}}{2} \approx 1.936$	Take the square root only at the end.

Problem 9

A closed cylindrical can must hold 16π cm³. Which radius and height use the least metal (the least total surface area), and what is that area?

$\pi r^2 h = 16\pi$, so $h = \frac{16}{r^2}$	The volume constraint, solved for the height.
$S(r) = 2\pi r^2 + 2\pi r h = 2\pi r^2 + \frac{32\pi}{r}$, for $r > 0$	Two ends plus the side; substitute h so that S depends on r alone.
$S'(r) = 4\pi r - \frac{32\pi}{r^2} = 0$, so $r^3 = 8$	Multiply by r^2 and divide by 4π .
$r = 2$, and $S''(r) = 4\pi + \frac{64\pi}{r^3} > 0$	A minimum, and the only critical point; $S \rightarrow \infty$ as $r \rightarrow 0$ and as $r \rightarrow \infty$, so it is global.

$h = \frac{16}{4} = 4$ and $S = 8\pi + 16\pi = 24\pi$	The height equals the diameter.
$r = 2$ cm , $h = 4$ cm , area $24\pi \approx 75.40$ cm²	For any volume the optimal can has $h = 2r$.

Problem 10

A model predicts the same number m for the four targets 1, 2, 6 and 7. Which m minimizes the squared error $E(m) = \sum_i (m - a_i)^2$, and what is the minimum error?

$E'(m) = \sum_i 2(m - a_i) = 2(4m - 16)$	Chain rule on each term; $\sum_i a_i = 16$.
$m = 4$	$E'(m) = 0$ gives $m = \frac{16}{4}$, the mean of the targets.
$E''(m) = 2 \cdot 4 = 8 > 0$	A minimum, and global: E is a parabola opening up. In general $E'' = 2n$.
$E(4) = 9 + 4 + 4 + 9 = 26$	The squared deviations from the mean.
$m = 4$, the mean; minimum squared error 26	The same calculation for any data shows the constant that minimizes squared error is the mean, which is why a regression's intercept fits the mean of the targets.

Where this goes wrong

1. Forgetting the endpoints

Setting the derivative to zero is the step that gets practiced, so it becomes the whole method.

$f(x) = x^3 - 3x$ on $[-2, 3]$, critical points $x = \pm 1$	Right so far: Problem 4, step 1.
“ $f(-1) = 2$ and $f(1) = -2$, so compare those.”	The habit that causes the mistake: only the critical points are evaluated.

The maximum is 2, at $x = -1$ ✗

That is only a local maximum. At the endpoint $x = 3$, $f(3) = 18$. On a closed interval the endpoints are always candidates, and here the maximum is at one, where $f'(3) = 24$, nowhere near 0.

2. Calling every flat point a maximum or minimum

$f'(x) = 0$ is the condition every extremum satisfies, and it is easy to read it the other way round.

$f(x) = x^4 - 4x^3$, $f'(x) = 4x^2(x - 3)$,
critical points 0 and 3

Right so far: Problem 2, step 2.

“ $f'(0) = 0$, so $x = 0$ is an extremum.”

The habit that causes the mistake: a zero derivative taken as enough.

$x = 0$ is a local maximum, with value
 $f(0) = 0$ ✗

f is decreasing on both sides of 0:
 $f(-1) = 5$ and $f(1) = -3$, and f' is
negative at -1 and at 1 . The tangent is
flat at 0 but the graph does not turn, like
 x^3 at 0. A critical point is a candidate;
the first-derivative test decides.

3. Reading an inconclusive test as “no extremum”

The second-derivative test usually gives an answer, so a zero feels like one.

$f(x) = x^4 - 4x^3 + 6x^2 - 4x$, $f'(x) =$
 $4(x - 1)^3$, critical point $x = 1$

Right so far: Problem 3, step 2.

“ $f''(1) = 0$, so $x = 1$ is not a maximum or
a minimum.”

The habit that causes the mistake:
 $f''(c) = 0$ treated as a third verdict,
“neither”.

f has no minimum ✗

$f''(1) = 0$ says nothing either way. The
first-derivative test does: $f'(0.9) =$
 -0.004 and $f'(1.1) = 0.004$, so f'
changes from $-$ to $+$ and $x = 1$ is a
minimum, with $f(1) = -1$. The same
happens for x^4 at 0.

4. Missing a critical point where f' is undefined

Critical points are usually found by solving $f'(x) = 0$, and a point where f' blows up is not a solution of anything.

$$f(x) = x^{2/3}(5 - x) \text{ on } [-1, 4], f'(x) = \frac{5(2 - x)}{3x^{1/3}}$$

Right so far: Problem 5, step 2.

“ $f'(x) = 0$ only at $x = 2$, so the candidates are $-1, 2$ and 4 .”

The habit that causes the mistake: only the zeros of the numerator are listed.

The minimum is $f(4) = 4^{2/3} \approx 2.520$ ✗

$f'(0)$ is undefined, so $x = 0$ is a critical point too, and $f(0) = 0$ is smaller. The graph has a cusp there: it falls into 0 from the left and rises out of it to the right. List the points where f' does not exist along with those where it is 0.

5. Answering with the location instead of the value

The derivative is solved for x , and the x that comes out feels like the answer.

$$f(x) = xe^{-x}, f'(x) = (1 - x)e^{-x},$$

maximum at $x = 1$

Right so far: Problem 6, step 3.

“The question asks for the maximum, and $x = 1$ is where it is.”

The habit that causes the mistake: stopping once the critical point is found.

The maximum value is 1 ✗

1 is the maximizer, where the maximum happens. The maximum value is $f(1) = \frac{1}{e} \approx 0.368$; f never reaches 1, since $xe^{-x} < 1$ for every x . Read the question again before writing the final line: Problem 7 asks for both, Problem 10 for both, this one for the value.